



Feb 2nd, 11:45 AM - 11:55 AM

Geovisualization of Ecological Data for Park Policy Support

Scott R. Abella

University of Nevada, Las Vegas

Haroon Stephen

University of Nevada, Las Vegas

Ross Guida

University of Nevada, Las Vegas

Scott R. Abella, Haroon Stephen, and Ross Guida, "Geovisualization of Ecological Data for Park Policy Support" (February 2, 2010).
2010 Annual Nevada NSF EPSCoR Climate Change Conference. Paper 39.
<http://digitalcommons.library.unlv.edu/epscor/2010/feb02/39>

This Event is brought to you for free and open access by the Conferences/Meetings (NNE) at University Libraries. It has been accepted for inclusion in 2010 Annual Nevada NSF EPSCoR Climate Change Conference by an authorized administrator of University Libraries. For more information, please contact marianne.buehler@unlv.edu.

Vegetation Change in the Newberry Mountains

Dr. Scott Abella


Dr. Haroon Stephen

Graduate student, Mr. Ross Guida

Policy and Outreach Component

Study Area

- Newberry Mountains, just NW of Laughlin
- Lake Mead NRA



Background

- Literature shows upward elevation shifts of biological species as a result of climate change
- Effects of climate change expected to accelerate in coming decades

Problem

- Concern both about species migrating out of parks and the potential inability of species to quickly adapt to the changing conditions within National Parks and other federal land boundaries

Research Questions

- What shifts in distributions of desert communities have occurred over the last 30 years across a 1,500 meter elevation gradient in the Newberry Mountains?
- Based on the changes in the last 30 years what potential future shifts may occur by the year 2040 under simulated climate change?

Datasets


- Baseline data: 1979 UNLV MS thesis by Jim Holland, park planner, Lake Mead NRA
 - 116 data points
- Points resurveyed in 2007-2008
- Datasets contain:
 - Slope
 - Aspect
 - Soil descriptions
 - Vegetation community data

Resurvey Field Photos



- Photos courtesy of Chris Roberts, LMNRA, Natural Resources Division

Data points in ArcGIS 9.3



Methods

- Maxent and Bioclim species distribution models will be used on 1979 data
- Determine model that most accurately portrays 2007-2008 species resurvey data
- Use model to project changes under simulated climate change 30 years from now
- ArcGIS used to create full databases and show changes over time
 - Shifts across the elevation gradient over the last 30 years

Project Goals

- Use the 30-year dataset to model past and potential desert community shifts
- Transfer findings to resource managers with NPS and other land management agencies
- Foster a meaningful two-way flow of data between researchers and stakeholders
- Produce original data and information which assists two-way flow of data into the future
- Produce geovisualizations in the new lab at UNLV
- Apply techniques on other study areas

Geovisualization

- 3D Visualization of mountains DEM overlaid with soil and vegetation maps
- Classification of satellite imagery using spectral signatures of known species
- Maps showing changes in density and distribution of vegetation
- Visualization for spatial characteristics of change (e.g., linking change at A to the change at B)
- Visualization of vegetation change in connection with climatic, ecological, meteorological, & hydrological variables
- Visualization of future projections/scenarios

References (see poster)

- Allen, C.D. and D.D. Breshears. 1998. Drought-induced shift of a forest-woodland ecotone: Rapid landscape response to climate variation. *Proceedings of the National Academy of Sciences* 94:9729-9733.
- Brown, J.H., T.J. Valone, and C.G. Curtin. 1997. Reorganization of an arid ecosystem in response to recent climate change. *Proceedings of the National Academy of Sciences* 94: 9729-9733.
- Daly, C. 2006. Guidelines for assessing the suitability of spatial climate data sets. *International Journal of Climatology* 26: 707-721.
- Dyer, J.M. 2002. A comparison of moisture scalars and water budget methods to assess vegetation-site relationships. *Physical Geography* 23: 245-258.
- Iverson, L.R., and A.M. Prasad. 1998. Predicting abundance of 80 tree species following climate change in the eastern United States. *Ecological Monographs* 68:465-485.
- Knapp, A.K., C. Beier, D.D. Briske, A.T. Classen, Y. Luo, M. Reichstein, M.D. Smith, S.D. Smith, J.E. Bell, P.A. Fay, J.L. Heisler, S.W. Leavitt, R. Sherry, B. Smith, and E. Weng. 2008. Consequences of more extreme precipitation regimes for terrestrial ecosystems. *BioScience* 58:811-821.
- Millar, C.I., and L.B. Brubaker. 2006. Climate change and paleoecology; new contexts for restoration ecology. Pp. 315-340 in Palmer, M., D. Falk, and J. Zedler (eds). *Restoration science*. Island Press, Washington, D.D.
- Penuelas, J., and M. Boada. 2003. A global change-induced biome shift in the Montseny Mountains (NE Spain). *Global Change Biology* 9:131-140.
- Phillips, S.J., R.P. Anderson, and R.E. Schapire. 2006. Maximum entropy modeling of species geographic distributions. *Ecological Modelling* 190: 231-259.
- Rice, K.J., and N.C. Emery. 2003. Managing microevolution: restoration in the face of global change. *Frontiers in Ecology and the Environment* 1:469-478.
- Sexton, J.P., and A.B. Grith. In preparation. Applying evolution to climate change management. Unpublished manuscript provided to Joshua Tree National Park, Twentyline Palms, CA.
- Walther, G., S. Beibner, and C.A. Burga. 2005. Trends in the upward shift of alpine plants. *Journal of Vegetation Science* 16: 541-548.
- Wardle, P., and M.C. Coleman. 1992. Evidence for rising upper limits of four native New Zealand forest trees. *New Zealand Journal of Botany* 30:303-314.